

Estimation of the Probability in the System of "Classifiers"

Alexsander Vardy

*The Department of Electrical and Computer Engineering
University of California, San Diego, USA, vardy@ece.ucsd.edu*

Oleg Kupervasser

*Department of Chemical Physics,
The Weizmann Institute of Science, Rehovot 76100, Israel, oleg_kup@yahoo.com*

Let that two types of recognition objects exist. Let that probabilities find these two types are known. Two variables that characterizes object exist and conditional probabilities to find one from two types of object exist as function every one variable. The best approximation for conditional probabilities as function of the two variables is necessary to find. The fuzzy logic usually used is not relevant. Cases of larger number of types and variables are considered.

PACS numbers 47.27.Gs, 47.27.Jv, 05.40.+j

I. INTRODUCTION

A. Statement of the problem.

Let A be a random variable, with values in set $0, 1$. Assume that the *a priori* probability $P(A) = P(A = 1)$ is known and denote it by θ . Let X_1, X_2 be two random variables, with values in some set, say $]-\infty; +\infty[$. We are given the following information: $X_1 = x_1$ and $X_2 = x_2$ (obtained though measurement). Furthermore, we have two system - "classifiers", which given x_1 and x_2 produce:

$$P(A = 1/X_1 = x_1) = P(A/x_1) \doteq \alpha \quad (1)$$

$$P(A = 1/X_2 = x_2) = P(A/x_2) \doteq \beta \quad (2)$$

We wish to estimate the probability $P(A = 1/X_1 = x_1, X_2 = x_2) = P(A/x_1, x_2)$ in terms of α, β and θ . More specifically we wish to find a function $\Gamma_{opt}(\alpha, \beta, \theta)$ which on the average is the best approximation for $P(A/x_1, x_2)$ in a sense to be defined explicitly in the sequel (see Fig 1.).

B. Importance of the problem

The problem is important for artificial intellect and can be widely used for image recognition.

II. NOTATION AND PRELIMINARIES

$\rho_{X_1, X_2}(x_1, x_2)$ - joint PDF(probability density function) of X_1 and X_2 . $\rho_{X_1, X_2/A}(x_1, x_2) \doteq h(x_1, x_2)$ - joint PDF of X_1 and X_2 , given $A = 1$. In terms of $h(x_1, x_2)$ and θ we may write $P(A/x_1, x_2)$ as follows:

$$P(A/x_1, x_2) = \frac{\theta h(x_1, x_2)}{\theta h(x_1, x_2) + (1 - \theta)\bar{h}(x_1, x_2)} \quad (3)$$

where $\bar{h}(x_1, x_2) \doteq \rho_{X_1, X_2/\bar{A}}(x_1, x_2)$ - joint PDF of X_1 and X_2 , given $A = 0$.

We have:

$$\rho_{X_1}(x_1) = \int_{-\infty}^{+\infty} \rho_{X_1, X_2}(x_1, x_2) dx_2 \quad (4)$$

$$\rho_{X_2}(x_2) = \int_{-\infty}^{+\infty} \rho_{X_1, X_2}(x_1, x_2) dx_1 \quad (5)$$

$$h_1(x_1) \doteq \rho_{X_1/A}(x_1) = \int_{-\infty}^{+\infty} h(x_1, x_2) dx_2 \quad (6)$$

$$h_2(x_2) \doteq \rho_{X_2/A}(x_2) = \int_{-\infty}^{+\infty} h(x_1, x_2) dx_1 \quad (7)$$

$$\bar{h}_1(x_1) \doteq \rho_{X_1/\bar{A}}(x_1) = \int_{-\infty}^{+\infty} \bar{h}(x_1, x_2) dx_2 \quad (8)$$

$$\bar{h}_2(x_2) \doteq \rho_{X_2/\bar{A}}(x_2) = \int_{-\infty}^{+\infty} \bar{h}(x_1, x_2) dx_1 \quad (9)$$

III. GENERIC FORM OF $P(A/X_1, X_2)$

Define the function $g(x_1, x_2)$ and $\bar{g}(x_1, x_2)$

$$g(x_1, x_2) \doteq \frac{h(x_1, x_2)}{h_1(x_1)h_2(x_2)} \quad (10)$$

$$\bar{g}(x_1, x_2) \doteq \frac{\bar{h}(x_1, x_2)}{\bar{h}_1(x_1)\bar{h}_2(x_2)} \quad (11)$$

Note that if X_1 and X_2 are conditionally independent, i.e.

$$h(x_1, x_2) = \rho_{X_1 X_2/A}(x_1, x_2) = \rho_{X_1/A}(x_1)\rho_{X_2/A}(x_2) = h_1(x_1)h_2(x_2) \quad (12)$$

then

$$g(x_1, x_2) = \bar{g}(x_1, x_2) = 1 \quad (13)$$

Define the following *monotonously nondecreasing* probability distribution functions

$$H_1(x_1) \doteq \int_{-\infty}^{x_1} h_1(z)dz \quad (14)$$

$$H_2(x_2) \doteq \int_{-\infty}^{x_2} h_2(z)dz \quad (15)$$

$$\bar{H}_1(x_1) \doteq \int_{-\infty}^{x_1} \bar{h}_1(z)dz \quad (16)$$

$$\bar{H}_2(x_2) \doteq \int_{-\infty}^{x_2} \bar{h}_2(z)dz \quad (17)$$

Note that since $H_1(x_1), H_2(x_2), \bar{H}_1(x_1)$ and $\bar{H}_2(x_2)$ are monotonous (At this point one could assume that $h_1(x_1), h_2(x_2), \bar{h}_1(x_1), \bar{h}_2(x_2) > 0$, so that $H_1(x_1), H_2(x_2), \bar{H}_1(x_1)$ and $\bar{H}_2(x_2)$ are monotonously increasing. This restriction will be shown to be superfluous in the sequel.) , there exist the inverse functions $H_1^{-1}(x_1), H_2^{-1}(x_2), \bar{H}_1^{-1}(x_1)$ and $\bar{H}_2^{-1}(x_2)$. We may therefore define:

$$J(a, b) \doteq g(H_1^{-1}(a), H_2^{-1}(b)) \quad (18)$$

$$\bar{J}(a, b) \doteq \bar{g}(\bar{H}_1^{-1}(a), \bar{H}_2^{-1}(b)) \quad (19)$$

For the sake of brevity we shall henceforth denote

$$J \doteq J(H_1(x_1), H_2(x_2)) = g(H_1^{-1}(H_1(x_1)), H_2^{-1}(H_2(x_2))) = g(x_1, x_2) \quad (20)$$

$$\bar{J} \doteq \bar{J}(\bar{H}_1(x_1), \bar{H}_2(x_2)) = \bar{g}(\bar{H}_1^{-1}(\bar{H}_1(x_1)), \bar{H}_2^{-1}(\bar{H}_2(x_2))) = \bar{g}(x_1, x_2) \quad (21)$$

By the definition

$$h(x_1, x_2) = Jh_1(x_1)h_2(x_2) \quad (22)$$

$$\bar{h}(x_1, x_2) = \bar{J}\bar{h}_1(x_1)\bar{h}_2(x_2) \quad (23)$$

We now have:

$$h_1(x_1) \doteq \rho_{X_1/A}(x_1) = \frac{\rho_{X_1}(x_1)P(A/x_1)}{P(A)} = \frac{\alpha\rho_{X_1}(x_1)}{\theta} . \quad (24)$$

$$h_2(x_2) \doteq \rho_{X_2/A}(x_2) = \frac{\rho_{X_2}(x_2)P(A/x_2)}{P(A)} = \frac{\beta\rho_{X_2}(x_2)}{\theta} . \quad (25)$$

$$\bar{h}_1(x_1) \doteq \rho_{X_1/\bar{A}}(x_1) = \frac{\rho_{X_1}(x_1)P(\bar{A}/x_1)}{P(\bar{A})} = \frac{(1-\alpha)\rho_{X_1}(x_1)}{1-\theta} \quad (26)$$

$$\bar{h}_2(x_2) \doteq \rho_{X_2/\bar{A}}(x_2) = \frac{\rho_{X_2}(x_2)P(\bar{A}/x_2)}{P(\bar{A})} = \frac{(1-\alpha)\rho_{X_2}(x_2)}{1-\theta} \quad (27)$$

Hence, from (22),(23)

$$h(x_1, x_2) = J \frac{\alpha\beta\rho_{X_1}(x_1)\rho_{X_2}(x_2)}{\theta^2} \quad (28)$$

$$\bar{h}(x_1, x_2) = \bar{J} \frac{(1-\alpha)(1-\beta)\rho_{X_1}(x_1)\rho_{X_2}(x_2)}{(1-\theta)^2} \quad (29)$$

Now from (3)

$$P(A/x_1, x_2) = \frac{\frac{J}{\theta}\alpha\beta\rho_{X_1}(x_1)\rho_{X_2}(x_2)}{\frac{J}{\theta}\alpha\beta\rho_{X_1}(x_1)\rho_{X_2}(x_2) + \frac{\bar{J}}{(1-\theta)}(1-\alpha)(1-\beta)\rho_{X_1}(x_1)\rho_{X_2}(x_2)} = \frac{\alpha\beta}{\alpha\beta + \frac{\bar{J}}{J}\frac{\theta}{1-\theta}(1-\alpha)(1-\beta)} \quad (30)$$

Note that in case of conditional independence $J = \bar{J} = 1$ and (30) becomes the exact solution $\Gamma(\alpha, \beta, \theta) = P(A/x_1, x_2)$.

IV. RESTRICTIONS ON THE FUNCTIONS $J(A, B)$ AND $\bar{J}(A, B)$

We have

$$h_1(x_1) = \int_{-\infty}^{+\infty} J(H_1(x_1), H_2(x_2))h_1(x_1)h_2(x_2)dx_2 . \quad (31)$$

Hence

$$1 = \int_{-\infty}^{+\infty} J(H_1(x_1), H_2(x_2)) h_2(x_2) dx_2 = \int_0^1 J(H_1(x_1), H_2(x_2)) dH_2(x_2) \quad (32)$$

Thus we have the following condition

$$\int_0^1 J(a, b) db = 1 \quad (33)$$

and analogously

$$\int_0^1 J(a, b) da = 1 \quad (34)$$

Similarly we obtain:

$$\begin{aligned} \int_0^1 \bar{J}(a, b) da &= 1 \\ \int_0^1 \bar{J}(a, b) db &= 1 \end{aligned} \quad (35)$$

Obviously

$$J(a, b), \bar{J}(a, b) \geq 0. \quad (36)$$

$$\int_0^1 \int_0^1 J(a, b) dadb = \int_0^1 \int_0^1 \bar{J}(a, b) dadb = 1. \quad (37)$$

The set of all the solutions of (33),(34),(35),(36),(37) together with (30) determines the set of all possible realizations of $P(A/x_1, x_2)$.

An example of a solution of (33),(34) and (36),(37).

Let $\rho(x)$ be a function such that $\rho(x) \geq 0$ and $\int_0^1 \rho(x) dx = 1$

Then

$$\begin{aligned} J(a, b) &= \rho(a - b) \quad , a \geq b \\ J(a, b) &= \rho(a - b + 1) \quad , a < b \end{aligned} \quad (38)$$

satisfies (33),(34) and (36),(37).

V. DEFINITION OF DISTANCE

We define the distance between the proposed approximation of $P(A/x_1, x_2)$, $\Gamma(\alpha, \beta, \theta)$ and the actual function $P(A/x_1, x_2)$ as follows:

$$\begin{aligned} ||\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2)|| &\doteq \\ \int \int_{-\infty}^{+\infty} \rho_{X_1 X_2}(x_1, x_2) & \\ [\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2)]^2 dx_1 dx_2 & \end{aligned} \quad (39)$$

Now we have from (22),(23) and (24),(25), (26),(27)

$$\begin{aligned} \rho_{X_1 X_2}(x_1, x_2) &= \theta h(x_1, x_2) + (1 - \theta) \bar{h}(x_1, x_2) = \\ \theta J h_1(x_1) h_2(x_2) &+ (1 - \theta) \bar{J} \bar{h}_1(x_1) \bar{h}_2(x_2) = \\ \left[\frac{J\alpha\beta}{\theta} + \frac{\bar{J}(1-\alpha)(1-\beta)}{(1-\theta)} \right] &\rho_{X_1}(x_1) \rho_{X_2}(x_2) \end{aligned} \quad (40)$$

$$\begin{aligned} ||\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2)|| & \\ = \int \int_{-\infty}^{+\infty} \rho_{X_1}(x_1) \rho_{X_2}(x_2) & \\ \left[\frac{J\alpha\beta}{\theta} + \frac{\bar{J}(1-\alpha)(1-\beta)}{(1-\theta)} \right] & \\ (\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2))^2 dx_1 dx_2 & \\ = \int_0^1 \int_0^1 \left[\frac{J\alpha\beta}{\theta} + \frac{\bar{J}(1-\alpha)(1-\beta)}{(1-\theta)} \right] & \\ (\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2))^2 dF_1(x_1) dF_2(x_2) & \end{aligned} \quad (41)$$

where

$$F_1(x_1) = \int_{-\infty}^{x_1} \rho_{X_1}(z) dz \quad (42)$$

$$F_2(x_2) = \int_{-\infty}^{x_2} \rho_{X_2}(z) dz \quad (43)$$

VI. RESTRAINTS FOR BASIC FUNCTIONS

We will consider in further all functions with arguments $1 \geq F_1, F_2 \geq 0$, but not x_1, x_2 . We have six function of F_1, F_2 , that define (41): $J, \bar{J}, H_1, H_2, \alpha, \beta$. Let us to write the other function by help these function and find restraints for there functions.

(i)

$$\alpha = P(A/x_1) = \theta h_1(x_1) / \rho_{X_1}(x_1) = \theta \frac{\frac{dH_1}{dx_1}}{\frac{dF_1}{dx_1}} = \theta \frac{dH_1}{dF_1} \quad (44)$$

By the same way

$$\beta = \theta \frac{dH_2}{dF_2} \quad (45)$$

Restraints for function H_1, H_2 are next:

$$\begin{aligned} H_1(1) &= H_2(1) = 1 \\ H_1(0) &= H_2(0) = 0 \end{aligned} \quad (46)$$

$$0 \leq \alpha = \theta \frac{dH_1}{dF_1}, \beta = \theta \frac{dH_2}{dF_2} \leq 1 \quad (47)$$

$$0 \leq \theta \leq 1 \quad (48)$$

(ii)

$$\begin{aligned}
\overline{H}_1(x_1) &= \int_{-\infty}^{x_1} \overline{h}_1(x_1) = \int_{-\infty}^{x_1} \frac{(1-\alpha)\rho_{X_1}(x_1)}{1-\theta} dx_1 = \\
&= \frac{1}{1-\theta} \int_{-\infty}^{x_1} -\frac{\theta}{1-\theta} \int_{-\infty}^{x_1} \frac{\alpha\rho_{X_1}(x_1)}{\theta} dx_1 \\
&= \frac{F_1}{1-\theta} - \frac{\theta}{1-\theta} H_1(x_1)
\end{aligned} \tag{49}$$

By the same way

$$\overline{H}_2(x_2) = \frac{F_2}{1-\theta} - \frac{\theta}{1-\theta} H_2(x_2) \tag{50}$$

(iii)

$$\begin{aligned}
&J(H_1(F_1), H_2(F_2)) : \\
&J(H_1(F_1), H_2(F_2)) \geq 0 \\
&\int_0^1 J(a, b) db = 1 \\
&\int_0^1 J(a, b) da = 1
\end{aligned} \tag{51}$$

$$\begin{aligned}
&\overline{J}(\overline{H}_1(F_1), \overline{H}_2(F_2)) : \\
&\overline{J}(\overline{H}_1(F_1), \overline{H}_2(F_2)) \geq 0 \\
&\int_0^1 \overline{J}(a, b) db = 1 \\
&\int_0^1 \overline{J}(a, b) da = 1
\end{aligned} \tag{52}$$

(iv)

$$P(A/x_1, x_2) = \frac{\frac{J\alpha\beta}{\theta}}{\frac{J\alpha\beta}{\theta} + \frac{\overline{J}(1-\alpha)(1-\beta)}{1-\theta}} \tag{53}$$

VII. OPTIMIZATION

We shall find the best approximation $\Gamma(\alpha, \beta, \theta)$ as follows

$$\min_{\Gamma(\alpha, \beta, \theta)} F[|\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2)|] \longrightarrow \Gamma(\alpha, \beta, \theta) \tag{54}$$

where the expected value $F[\dots]$ is taken with respect to the joint PDF of possible realizations of: $J, \overline{J}, \alpha, \beta, H_1, H_2$ for given F_1 and F_2 .

For the sake of brevity we denote:

$$C \doteq \frac{J\alpha\beta}{\theta} + \frac{\overline{J}(1-\alpha)(1-\beta)}{(1-\theta)} \tag{55}$$

$$D \doteq \frac{J\alpha\beta}{\theta} \tag{56}$$

Then from (53) and (41)

$$\begin{aligned}
&||\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2)|| = \\
&\int_0^1 \int_0^1 C(\Gamma(\alpha, \beta, \theta) - D/C)^2 dF_1 dF_2 = \\
&\int_0^1 \int_0^1 dF_1 dF_2 [\frac{D^2}{C} + \Gamma^2(\alpha, \beta, \theta)C - 2\Gamma(\alpha, \beta, \theta)D]
\end{aligned} \tag{57}$$

Thus

$$\begin{aligned}
&\min_{\Gamma(\alpha, \beta, \theta)} F[|\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2)|] = \\
&\min_{\Gamma(\alpha, \beta, \theta)} F[\int_0^1 \int_0^1 dF_1 dF_2 \\
&[\frac{D^2}{C} + \Gamma^2(\alpha, \beta, \theta)C - 2\Gamma(\alpha, \beta, \theta)D]] = \\
&\min_{\Gamma(\alpha, \beta, \theta)} F[\int_0^1 \int_0^1 dF_1 dF_2 [\frac{D^2}{C}]] + \\
&\min_{\Gamma(\alpha, \beta, \theta)} F[\int_0^1 \int_0^1 dF_1 dF_2 \\
&[\Gamma^2(\alpha, \beta, \theta)C - 2\Gamma(\alpha, \beta, \theta)D]] = \\
&Const + \min_{\Gamma(\alpha, \beta, \theta)} F[\int_0^1 \int_0^1 dF_1 dF_2 \\
&[\Gamma^2(\alpha, \beta, \theta)C - 2\Gamma(\alpha, \beta, \theta)D]]
\end{aligned} \tag{58}$$

It remains to calculate the expected value in (58).

We have by obvious assumptions

$$\begin{aligned}
&\rho_{J, \overline{J}, \alpha, \beta, H_1, H_2 / F_1, F_2}(J, \overline{J}, \alpha, \beta, H_1, H_2 / F_1, F_2) = \\
&\rho_{J / H_1, H_2}(J / H_1, H_2) \rho_{\overline{J} / \overline{H}_1, \overline{H}_2}(\overline{J} / \overline{H}_1, \overline{H}_2) \\
&\rho_{\alpha / F_1}(\alpha / F_1) \rho_{H_1 / \alpha, F_1}(H_1 / \alpha, F_1) \\
&\rho_{\beta / F_2}(\beta / F_2) \rho_{H_2 / \beta, F_2}(H_2 / \beta, F_2)
\end{aligned} \tag{59}$$

A. Lemma 1

$$F[J(a, b)] = \int_0^{+\infty} \rho_{J(a, b) / a, b}(J(a, b) / a, b) J(a, b) dJ = 1 \tag{60}$$

$$F[\overline{J}(a, b)] = \int_0^{+\infty} \rho_{\overline{J}(a, b) / a, b}(\overline{J}(a, b) / a, b) \overline{J}(a, b) d\overline{J} = 1 \tag{61}$$

Proof:

Let us consider function: $\rho_{J(a, b) / a, b}$ Function $J(a, b)$ is defined on the square $0 \leq a, b \leq 1$ Let us make sampling of function J on this square by its dividing on small squares (i, j) and define value of the function J_{ij} on every square i, j . Restraints for function J (***) can be written

$$J_{ij} \geq 0 \tag{62}$$

$$\frac{1}{N} \sum_{i=1}^N J_{ij} = 1 \quad (63)$$

$$\frac{1}{N} \sum_{j=1}^N J_{ij} = 1 \quad (64)$$

here $i = 1, \dots, N, j = 1, \dots, N$

All matrixes (J_{ij}) that satisfy these conditions are equal probability. Let us define probability density function

$$\rho(J_{11}, \dots, J_{ij}, \dots, J_{NN}) \quad (65)$$

From symmetry conditions that define this function (ρ) with respect to transpositions lines and columns in matrix (J_{ij}) we can conclude that this function (ρ) also doesn't transform with respect to such transpositions.

Let us consider function $\rho_{u/ij}(u/ij)$ which is a discrete version of the function $\rho_{J(a,b)/a,b}(J(a,b)/a,b)$:

$$\rho_{u/ij}(u/ij) = \int \dots \int_0^{+\infty} \rho(J_{11}, \dots, J_{nk}, \dots, J_{ij} = u, \dots, J_{NN}) \prod_{(lm) \neq (ij)} dJ_{lm} \quad (66)$$

Let us transpose lines and columns (J_{ij}) by such way that element J_{ij} will be replaced by element J_{nk} , the function $\rho(J_{11}, \dots)$ will not be transform after it. So from previous equation we obtain

$$\rho_{u/ij}(u/ij) = \int \dots \int_0^{+\infty} \rho(J_{11}, \dots, J_{nk} = u, \dots, J_{ij}, \dots, J_{NN}) \prod_{(lm) \neq (nk)} dJ_{lm} = \rho_{u/nk}(u/nk) \quad (67)$$

From this equation we can conclude that $\rho_{u/ij}(u/ij)$ doesn't depend on ij so $\rho_{J/ab}(J/ab)$ doesn't depend on ab and

$$\rho_{J/ab}(J/ab) = \rho_J(J) \quad (68)$$

And

$$F[J(a,b)] = \int_0^{+\infty} \rho_J(J) J dJ = Const \quad (69)$$

From

$$\int_0^1 \int_0^1 J(a,b) dadb = 1 \quad (70)$$

we can conclude that

$$\int_0^1 \int_0^1 F[J(a,b)] dadb = 1 \quad (71)$$

So we can obtain that $Const = 1$ in Eq.(69).

B. Lemma 2

Probability distribution functions α and β doesn't depend on F_1 and F_2 .

$$\rho_{\alpha/F_1}(\alpha/F_1) = \rho_\alpha(\alpha) \quad (72)$$

$$\rho_{\beta/F_2}(\beta/F_2) = \rho_\beta(\beta) \quad (73)$$

Proof:

Let us make sampling of function $\alpha(F_1)$ by dividing of domain of this function $F_1, [0, 1]$ on intervals of $1/N, N \gg 1$ Then restriction conditions for $\alpha_k, k = 1, \dots, N$:

$$0 \leq \alpha_k \leq 1 \quad (74)$$

$$\frac{1}{N} \sum_{k=1}^N \alpha_k = \int_0^1 \theta \frac{dH_1(F_1)}{dF_1} dF_1 = \theta \quad (75)$$

All columns (α_k) that satisfy by these conditions are equal probability. Let us consider respective function $\rho(\alpha_1, \dots, \alpha_k, \dots, \alpha_l, \dots, \alpha_N)$. From symmetry conditions that define this function with respect to transpositions $\alpha_k \rightarrow \alpha_l$ function $\rho(\alpha_1, \dots, \alpha_k, \dots, \alpha_l, \dots, \alpha_N)$ also doesn't transform with respect to such transpositions. So we can write

$$\begin{aligned} \rho_k(u) &= \int_0^1 \rho(\alpha_1, \dots, \alpha_k = u, \dots, \alpha_l, \dots, \alpha_N) \prod_{n \neq k} d\alpha_n = \\ &= \int_0^1 \rho(\alpha_1, \dots, \alpha_k, \dots, \alpha_l = u, \dots, \alpha_N) \prod_{n \neq l} d\alpha_n = \\ &= \rho_l(u) \end{aligned} \quad (76)$$

From this equation we can conclude that function $\rho_{\alpha/F_1}(\alpha/F_1)$ doesn't depend on F_1 .

$$\rho_{\alpha/F_1}(\alpha/F_1) = \rho_\alpha(\alpha) \quad (77)$$

From (59) we obtain

$$\begin{aligned} F[\Gamma^2(\alpha, \beta, \theta)C - 2\Gamma(\alpha, \beta, \theta)D] &= \\ &= \int_0^1 \int_0^1 \rho_\alpha(\alpha) \rho_\beta(\beta) d\alpha d\beta \\ &= \int_0^1 \int_0^1 \rho_{H_1/\alpha, F_1}(H_1/\alpha, F_1) \rho_{H_2/\beta, F_2}(H_2/\beta, F_2) dH_1 dH_2 \\ &= \int_0^\infty \int_0^\infty \rho_J(J) \rho_{\bar{J}}(\bar{J}) [\Gamma^2(\alpha, \beta, \theta) [\frac{J\alpha\beta}{\theta} + \frac{\bar{J}(1-\alpha)(1-\beta)}{1-\theta}] \\ &\quad - 2\Gamma(\alpha, \beta, \theta) \frac{J\alpha\beta}{\theta}] dJ d\bar{J} \\ &= \int_0^1 \int_0^1 \rho_\alpha(\alpha) \rho_\beta(\beta) d\alpha d\beta \\ &= [\Gamma^2(\alpha, \beta, \theta) [\frac{F[J]\alpha\beta}{\theta} + \frac{F[\bar{J}](1-\alpha)(1-\beta)}{1-\theta}] \\ &\quad - 2\Gamma(\alpha, \beta, \theta) \frac{F[J]\alpha\beta}{\theta}] \end{aligned} \quad (78)$$

Let us define

$$\overline{C} = \frac{\alpha\beta}{\theta} + \frac{(1-\alpha)(1-\beta)}{1-\theta} \quad (79)$$

$$\overline{D} = \frac{\alpha\beta}{\theta} \quad (80)$$

By Lemma 1 , $F[J] = F[\overline{J}] = 1$. Hence

$$F[\Gamma^2(\alpha, \beta, \theta)C - 2\Gamma(\alpha, \beta, \theta)D] = \int_0^1 \int_0^1 [\Gamma^2(\alpha, \beta, \theta)\overline{C} - 2\Gamma(\alpha, \beta, \theta)\overline{D}] \rho_\alpha(\alpha) \rho_\beta(\beta) d\alpha d\beta \quad (81)$$

It remain to find

$$\begin{aligned} & \min_{\Gamma(\alpha, \beta, \theta)} \int_0^1 \int_0^1 dF_1 dF_2 \\ & \int_0^1 \int_0^1 d\alpha d\beta \rho_\alpha(\alpha) \rho_\beta(\beta) \\ & [\Gamma^2(\alpha, \beta, \theta)\overline{C} - 2\Gamma(\alpha, \beta, \theta)\overline{D}] \end{aligned} \quad (82)$$

Since

$$\rho_\alpha(\alpha) \rho_\beta(\beta) \geq 0 \quad (83)$$

if the expression in square brackets is minimized at each point then the whole integral in (82) is minimized. Thus we may proceed as follows

$$\frac{\partial}{\partial \Gamma} [\Gamma^2(\alpha, \beta, \theta)\overline{C} - 2\Gamma(\alpha, \beta, \theta)\overline{D}] = 2\Gamma(\alpha, \beta, \theta)\overline{C} - 2\overline{D} = 0 \quad (84)$$

Hence the optimum $\Gamma(\alpha, \beta, \theta)$ is given by

$$\Gamma_{opt}(\alpha, \beta, \theta) = \frac{\overline{D}}{\overline{C}} = \frac{\frac{\alpha\beta}{\theta}}{\frac{\alpha\beta}{\theta} + \frac{(1-\alpha)(1-\beta)}{1-\theta}} \quad (85)$$

VIII. MEAN DISTANCE BEETWEEN THE PROPOSED APPROXIMATION OF $P(A/X_1, X_2)$, - $\Gamma(\alpha, \beta, \theta)$ AND THE ACTUAL FUNCTION $P(A/X_1, X_2)$

The mean distance from (57) is

$$\begin{aligned} DIS &= F[|\Gamma(\alpha, \beta, \theta) - P(A/x_1, x_2)|] = \\ & \int_0^1 \int_0^1 \rho_\alpha(\alpha) \rho_\beta(\beta) d\alpha d\beta \\ & [\Gamma^2(\alpha, \beta, \theta)\overline{C} - 2\Gamma(\alpha, \beta, \theta)\overline{D}] \\ & + Const \end{aligned} \quad (86)$$

where $Const$ in this equation is defined by

$Const =$

$$F\left[\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \rho_{X_1, X_2}(x_1, x_2) [P(A/x_1, x_2)]^2 dx_1 dx_2\right] \quad (87)$$

From this equation we can find boundaries of the $Const$. From $0 \leq P(A/x_1, x_2) \leq 1$ we can conclude

$$\begin{aligned} Const &\leq \\ F\left[\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \rho_{X_1, X_2}(x_1, x_2) P(A/x_1, x_2) dx_1 dx_2\right] \\ &= F[\theta] = \theta \end{aligned} \quad (88)$$

The second condition is

$$\begin{aligned} 0 &\leq F\left[\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \rho_{X_1, X_2}(x_1, x_2) \right. \\ & [P(A/x_1, x_2) - \theta]^2 dx_1 dx_2] = \\ & F\left[\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \rho_{X_1, X_2}(x_1, x_2) \right. \\ & [P(A/x_1, x_2)^2 + \theta^2 - 2P(A/x_1, x_2)\theta] dx_1 dx_2] = \\ & F\left[\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \rho_{X_1, X_2}(x_1, x_2) [P(A/x_1, x_2)]^2 dx_1 dx_2\right] \\ & - \theta^2 \end{aligned} \quad (89)$$

So from this two equations we can conclude

$$\theta^2 \leq Const \leq \theta \quad (90)$$

By next step we would like find function $\rho_\alpha(\alpha)$ ($\rho_\beta(\beta)$) in equation for DIS .

Restrictions for function $\alpha(F_1)$, $0 \leq F_1 \leq 1$ are next:

(i)

$$\int_0^1 \alpha(F_1) dF_1 = \theta \quad (91)$$

(ii)

$$0 \leq \alpha(F_1) \leq 1 \quad (92)$$

In discrete form (for $N \rightarrow \infty$) we can rewrite $\alpha_{set} = \{\alpha_1, \alpha_2, \dots, \alpha_N\}$

(i)

$$\frac{1}{N} \sum_{i=1}^N \alpha_i = \theta \quad (93)$$

(ii)

$$0 \leq \alpha_i \leq 1, i = 1, 2, \dots, N \quad (94)$$

Let us define function $U(\alpha_{set})$ by next way

$$\begin{aligned} U(\alpha_{set}) &= \sum_{i=1}^N \alpha_i \quad \text{for } 0 \leq \alpha_i \leq 1, i = 1, 2, \dots, N \\ U(\alpha_{set}) &= +\infty \quad \text{otherwise} \end{aligned} \quad (95)$$

$$U(\alpha_{set}) = \sum_{i=1}^N U_i(\alpha_i) \quad (96)$$

$$\begin{aligned} U_i(\alpha_i) &= \alpha_i \quad \text{for } 0 \leq \alpha_i \leq 1 \\ U_i(\alpha_i) &= +\infty \quad \text{otherwise} \end{aligned} \quad (97)$$

Then function that satisfy equal probability distribution with considering restrictions (i),(ii) is

$$\rho_{\alpha_{set}}(\alpha_{set}) = \frac{1}{C} \delta(U(\alpha_{set}) - N\theta) \quad (98)$$

where δ - delta-function of Dirac.

Constant C define by

$$\int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \rho_{\alpha_{set}}(\alpha_{set}) d\alpha_1 \dots d\alpha_N = 1 \quad (99)$$

It can be proved (see each course of "Statistical mechanics"; transform from microcanonical to canonical distribution) that for $N \mapsto \infty$ distribution (98) is equal to next distribution:

$$\rho_{\alpha_{set}}(\alpha_{set}) = \frac{1}{Z} e^{-KU(\alpha_{set})} \quad (100)$$

where Z and K can be found from equations

$$\int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \rho_{\alpha_{set}}(\alpha_{set}) d\alpha_1 \dots d\alpha_N = 1 \quad (101)$$

$$\int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} U(\alpha_{set}) \rho_{\alpha_{set}}(\alpha_{set}) d\alpha_1 \dots d\alpha_N = N\theta \quad (102)$$

Quest function $\rho_\alpha(\alpha)$ can be find by

$$\begin{aligned} \rho_\alpha(\alpha) &= \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \rho_{\alpha_{set}}(\alpha_1, \dots, \alpha_j = \alpha, \dots, \alpha_N) \\ &\prod_{i=1, i \neq j}^N d\alpha_i = \\ &\frac{1}{D} e^{-KU_j(\alpha_j = \alpha)} \end{aligned} \quad (103)$$

where

$$D^N = Z \quad (104)$$

From Eqs.(101),(102) we can find

$$\frac{1}{Z} = \left(\frac{K}{1 - e^{-K}} \right)^N \quad (105)$$

$$\theta = \Lambda(K) \quad (106)$$

where $\Lambda(K)$ is decreasing function

$$\begin{aligned} \Lambda(K) &= 1 \quad \text{for } K = -\infty \\ \Lambda(K) &= 0 \quad \text{for } K = +\infty \\ \Lambda(K) &= 1/2 \quad \text{for } K = 0 \\ \Lambda(K) &= \frac{1}{K} - \frac{1}{e^K - 1} \quad \text{otherwise} \end{aligned} \quad (107)$$

If K is root of Eq (refpor6) we can write from Eqs.(103),(104),(105),(106) for function $\rho_\alpha(\alpha)$:

$$\begin{aligned} \text{For } K &= 0 \\ \rho_\alpha(\alpha) &= 1 \quad \text{for } 0 \leq \alpha \leq 1 \\ \rho_\alpha(\alpha) &= 0 \quad \text{otherwise} \\ \text{For } K &= +\infty \\ \rho_\alpha(\alpha) &= 2\delta(\alpha) \quad 0 \leq \alpha \leq 1 \\ \rho_\alpha(\alpha) &= 0 \quad \text{otherwise} \\ \text{For } K &= -\infty \\ \rho_\alpha(\alpha) &= 2\delta(\alpha - 1) \quad 0 \leq \alpha \leq 1 \\ \rho_\alpha(\alpha) &= 0 \quad \text{otherwise} \\ \text{For } \text{otherwise } K \\ \rho_\alpha(\alpha) &= \frac{1}{D} e^{-K\alpha} \quad 0 \leq \alpha \leq 1 \\ \rho_\alpha(\alpha) &= 0 \quad \text{otherwise} \end{aligned} \quad (108)$$

where $2 \int_0^1 \delta(\alpha - 1) = 2 \int_0^1 \delta(\alpha) = 1$ and

$$\frac{1}{D} = \frac{K}{1 - e^{-K}} \quad (109)$$

IX. THE CASE OF MORE THEN TWO VARIABLES A AND X

Let A be a random variable, with values in set $0, 1, \dots, L$. Assume that the *a priori* probability $P(A = i)$ is known and denote it by θ_i , here $i = 1, \dots, L$. Let $X_1, \dots, X_K$ be K random variables, with values in some set, say $] - \infty; +\infty[$. We are given the following information: $X_1 = x_1, \dots, X_K = x_K$ (obtained though measurement). Furthermore, we have two system - "classifiers", which given $x_1, \dots, x_K$ produce:

$$P(A = i / X_j = x_j) \doteq \alpha_{ij} \quad (110)$$

We wish to estimate the probability $P(A = i / X_1 = x_1, \dots, X_K = x_K)$ in terms of α_{ij} and θ_i . More specifically we wish to find a function $\Gamma_{opt,M}(\alpha_{ij}, \theta_i)$ which on the average is the best approximation for $P(A = M / x_1, \dots, x_K)$. By the same way that in case of two variables we can find that the $\Gamma_{opt,M}(\alpha_{ij}, \theta_i)$ defined by equation

$$\Gamma_{opt,M}(\alpha_{ij}, \theta_i) = \frac{(\prod_{j=1}^K \alpha_{Mj}) / \theta_M^{K-1}}{\sum_{i=1}^L (\prod_{j=1}^K \alpha_{ij}) / \theta_i^{K-1}} \quad (111)$$

We have evidential restraints for α_{ij}, θ_i

$$\begin{aligned} 0 &\leq \alpha_{ij} \leq 1 \\ \sum_{i=1}^L \alpha_{ij} &= 1 \end{aligned} \quad (112)$$

$$\begin{aligned} 0 &\leq \theta_i \leq 1 \\ \sum_{i=1}^L \theta_i &= 1 \end{aligned} \quad (113)$$

-
- [1] Landau L.D., Lifshits E.M. *Statisticheskaja fizika*, ("Nauka", Moskva(1976))
[2] Handbook, *Iskustvennyj intellekt*, Spravochnik, Pospelov D.A., ("Radio i svjaz" Moskva 1990)
[3] Ventsel' E.S., *Teorija verojatnostej*, ("Nauka", Moskva(1969))

Figures Legends

Fig.1: Function $\Gamma(\alpha, \beta, \theta) : [0, 1]^3 \mapsto [0, 1]$




Fig 1

FIG. 1. The System of "Classifiers".
Function $\Gamma(\alpha, \beta, \theta) : [0, 1]^3 \mapsto [0, 1]$